



HEDIS - Effectiveness of Preventive Care

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2009 through Dec 31, 2009

Location	Provider	Adult BMI	Breast Cancer Screening	Cervical Cancer Screening	Chlamydia Screening	Childhood Immunizations	Colorectal Cancer Screening	Lead Screening in Children	Child BMI
SETMA 1	Aziz	95.1%	65.2%	0.0%	--	--	97.7%	--	--
	Duncan	94.7%	71.5%	11.2%	--	--	98.7%	--	--
	Henderson	98.5%	74.0%	15.7%	--	--	97.3%	--	--
	Murphy	96.9%	57.8%	2.4%	--	--	96.0%	--	--
	Thomas	98.5%	61.5%	0.0%	--	--	97.8%	--	--
SETMA 1 Totals:		96.4%	68.3%	8.2%	--	--	97.6%	--	--
SETMA 2	Anthony	98.4%	78.9%	0.9%	--	--	98.0%	--	--
	Anwar	99.5%	83.5%	0.0%	--	--	99.1%	--	--
	Cricchio, M	100.0%	73.2%	0.0%	--	--	98.9%	--	--
	Holly	100.0%	82.9%	0.0%	--	--	100.0%	--	--
	Leifeste	99.4%	83.7%	3.8%	--	--	98.3%	--	--
	Wheeler	99.2%	80.4%	17.9%	--	--	97.8%	--	--
SETMA 2 Totals:		99.4%	80.6%	5.7%	--	--	98.6%	--	--
SETMA West	Curry	99.6%	72.5%	16.7%	--	--	99.6%	--	--
	Halbert	97.1%	73.7%	1.9%	--	--	98.2%	--	--
	Horn	99.5%	84.1%	9.4%	--	--	98.5%	--	--
	Satterwhite	99.6%	82.3%	9.0%	--	--	99.3%	--	--
	Vardiman	96.4%	54.7%	0.0%	--	--	96.9%	--	--
	Young	100.0%	74.5%	3.7%	--	--	95.9%	--	--
SETMA West Totals:		98.9%	78.2%	8.0%	--	--	98.4%	--	--
SETMA Totals:		98.3%	75.9%	7.3%	--	--	98.2%	--	--

Adult Body Mass Assessment - Calculation of Body Mass Index (BMI) for patients 18 to 74 years of age.

Breast Cancer Screening - Yearly mammography screening for women 40 to 69 years of age.

Cervical Cancer Screening - Cervical cancer screening for women 21 to 64 years of age at least every two years.

Chlamydia Screening - Sexually active females between the ages of 16 and 24 should have a Chlamydia screening at least yearly.



HEDIS - Effectiveness of Preventive Care

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2009 through Dec 31, 2009

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Colorectal Cancer Screening - Patients 50 to 80 years of age should have at least one of the following: a colonoscopy within the last nine years, a flexible sigmoidoscopy within the last four years, a double contrast barium enema within the last four years or a fecal occult blood test within the last year.

Lead Screening in Children - Children should have a serum lead screening by the time of their second birthday.

Childhood Body Mass Assessment - Children 2 to 18 years of age should have the Body Mass Index (BMI) percentile documented.