



NQF - Care for Older Adults

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2009 Through Dec 30, 2009

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA 1	Aziz	72.6%	63.3%	59.2%	13.1%
	Duncan	95.0%	73.0%	54.3%	20.5%
	Henderson	80.3%	60.3%	62.4%	42.0%
	Murphy	78.1%	47.0%	57.9%	18.2%
	Thomas	42.0%	49.3%	95.2%	54.1%
SETMA 1 Totals:		76.8%	58.2%	61.6%	24.6%
SETMA 2	Anthony	98.7%	76.3%	67.3%	37.8%
	Anwar	99.7%	82.0%	69.1%	36.7%
	Colbert	--	--	--	--
	Cricchio, M	99.2%	71.1%	58.8%	35.7%
	Holly	98.8%	79.3%	76.1%	42.5%
	Leifeste	92.1%	84.2%	60.3%	33.4%
	Wheeler	99.6%	65.8%	64.4%	43.4%
SETMA 2 Totals:		98.2%	77.2%	66.7%	37.8%
SETMA West	Anthony	--	--	--	--
	Curry	98.5%	44.4%	58.1%	28.5%
	Halbert	65.3%	41.6%	57.2%	35.3%
	Horn	72.9%	68.6%	77.8%	49.9%
	Satterwhite	89.3%	55.9%	51.6%	38.7%
	Vardiman	95.6%	53.6%	60.7%	27.2%
	Young	98.5%	39.0%	71.4%	39.6%
SETMA West Totals:		82.2%	49.5%	61.6%	36.3%
SETMA Totals:		86.3%	61.7%	63.5%	32.8%

Counseling For Physical Activity - Patients 65 and older should be counseled to engage in physical activity.

Colorectal Cancer Screening - Patients 50 to 80 years of age should have at least one of the following: a colonoscopy within the last nine years, a flexible sigmoidoscopy within the last four years, a double contrast barium enema within the last four years or a fecal occult blood test within the last year.



NQF - Care for Older Adults

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2009 Through Dec 30, 2009

Fall Risk Assessment - Patients 65 and older should have a fall risk assessment completed at least yearly.

Urinary Incontinence Assessment - Patients 65 and older should have urinary incontinence assessed at least every six months.