



NQF - Medication Measures - Persistent Medications

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2009 through Dec 31, 2009

Annual Monitoring for Patients on								
Location	Provider	Digoxin	Diuretics	ACE or ARB	Pheno-barbital	Carba-mazepine	Phenytoin	Valproic
SETMA 1	Aziz	94.3%	93.7%	94.2%	83.3%	27.3%	71.0%	22.9%
	Duncan	89.6%	88.1%	89.2%	100.0%	40.0%	65.2%	6.7%
	Henderson	96.1%	91.8%	92.3%	68.4%	36.4%	81.2%	25.0%
	Murphy	94.3%	94.4%	94.3%	70.0%	75.0%	73.5%	21.1%
	Thomas	86.8%	78.8%	79.1%	60.0%	0.0%	80.0%	41.7%
SETMA 1 Totals:		93.6%	91.0%	91.5%	75.0%	34.5%	73.4%	23.1%
SETMA 2	Anthony	93.2%	93.3%	93.4%	77.8%	28.6%	97.6%	53.3%
	Anwar	94.3%	93.9%	93.5%	92.6%	54.5%	83.1%	48.1%
	Colbert	--	100.0%	100.0%	--	--	--	0.0%
	Cricchio, M	91.3%	90.8%	91.2%	75.0%	30.0%	83.7%	18.8%
	Holly	96.7%	93.4%	93.3%	70.0%	12.5%	77.1%	23.1%
	Leifeste	92.5%	89.2%	89.7%	100.0%	16.7%	73.9%	40.9%
	Wheeler	81.2%	93.1%	92.8%	100.0%	33.3%	80.0%	50.0%
SETMA 2 Totals:		92.9%	92.4%	92.3%	86.2%	29.6%	83.3%	39.2%
SETMA West	Curry	86.5%	72.9%	73.5%	71.4%	0.0%	52.9%	23.1%
	Halbert	87.8%	74.2%	75.2%	35.0%	22.2%	35.6%	13.5%
	Horn	60.5%	64.8%	65.7%	66.7%	20.0%	53.8%	0.0%
	Satterwhite	89.1%	75.6%	76.4%	85.7%	11.1%	50.0%	1.8%
	Vardiman	67.5%	71.0%	72.8%	--	0.0%	62.5%	0.0%
	Young	79.5%	73.6%	73.4%	0.0%	30.0%	47.4%	9.1%
SETMA West Totals:		80.8%	72.2%	73.0%	45.5%	15.4%	45.1%	7.8%
SETMA Totals:		90.0%	85.6%	86.2%	71.9%	26.8%	70.5%	20.0%

Annual Monitoring for Patients on Persistent Medications

Patients with an active prescription of digoxin should have a serum potassium and a serum creatinine test or a serum potassium and a blood urea nitrogen test at least yearly.
 Patients with an active prescription of a diuretic should have a serum potassium and a serum creatinine test or a serum potassium and a blood urea nitrogen test at least yearly.
 Patients with an active prescription of an ACE or ARB should have a serum potassium and a serum creatinine test or a serum potassium and a blood urea nitrogen test at least yearly.
 Patients with an active prescription for an anticonvulsant should have a corresponding drug level checked at least yearly.