

HEDIS - Effectiveness of Chronic Care - Persistent Medications (MPM)

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

		Annual Monitoring for Patients on						
Location	Provider	Digoxin	Diuretics	ACE or ARB	Pheno- barbital	Carba- mazepine	Phenytoin	Valproic
SETMA 1	Anthony							
	Aziz	94.1%	96.2%	96.1%	76.7%	21.4%	84.5%	52.0%
	Duncan	97.1%	88.9%	90.7%		66.7%	84.2%	14.3%
	Henderson	98.7%	90.9%	91.0%	66.7%	50.0%	87.5%	53.7%
	Murphy	99.0%	96.0%	96.1%	88.9%	33.3%	58.0%	35.0%
	Thomas	90.8%	85.9%	86.3%	100.0%	0.0%	86.7%	44.7%
SETMA 1 Totals:		95.9%	92.3%	92.8%	80.0%	30.2%	78.3%	42.8%
SETMA 2	Anthony	95.8%	94.0%	95.2%	85.7%	33.3%	86.0%	69.0%
	Anwar	98.8%	95.8%	96.0%	82.6%	46.2%	81.2%	56.1%
	Colbert		75.0%	75.0%	0.0%		0.0%	
	Cricchio, A		100.0%					
	Cricchio, M	90.7%	93.8%	93.9%	100.0%	20.0%	65.0%	21.7%
	Holly	100.0%	96.5%	96.0%	100.0%	50.0%	84.2%	0.0%
	Leifeste	92.9%	91.2%	91.8%	100.0%	0.0%	77.3%	21.7%
	Wheeler	92.7%	92.5%	93.7%	90.9%	42.9%	91.7%	48.0%
SETMA 2 Totals:		96.1%	93.9%	94.4%	84.8%	31.9%	79.1%	44.5%
SETMA West	Curry	92.7%	84.0%	85.4%		0.0%	76.9%	21.7%
	Deiparine	75.4%	73.7%	73.6%	68.2%	40.9%	70.6%	42.9%
	Halbert	85.0%	79.6%	79.5%	33.3%	17.6%	51.7%	20.4%
	Horn	92.9%	83.4%	84.7%	0.0%	33.3%	82.1%	15.4%
	Qureshi	83.3%	78.1%	78.9%			100.0%	0.0%
	Satterwhite	95.2%	83.3%	84.5%	100.0%	20.0%	92.3%	2.6%
	Vardiman	83.3%	78.6%	79.4%	100.0%	50.0%	83.3%	25.0%
	Young	93.3%	75.1%	74.4%		0.0%	46.2%	38.5%
SETMA West Totals:		86.8%	80.0%	80.5%	56.8%	27.0%	72.0%	21.8%
SETMA Totals:		93.2%	88.8%	89.3%	75.0%	29.4%	76.8%	34.5%



HEDIS - Effectiveness of Chronic Care - Persistent Medications (MPM)

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Annual Monitoring for Patients on Persistent Medications

Patients with an active prescription of digoxin should have a serum potassium and a serum creatinine test or a serum potassium and a blood urea nitrogen test at least yearly. Patients with an active prescription of a diuretic should have a serum potassium and a serum creatinine test or a serum potassium and a blood urea nitrogen test at least yearly. Patients with an active prescription of an ACE or ARB should have a serum potassium and a serum creatinine test or a serum potassium and a blood urea nitrogen test at least yearly. Patients with an active prescription for an anticonvulsant should have a corresponding drug level checked at least yearly.

Nov 28, 2011 11:56:46 AM 2 of 2 NGHE060