

HEDIS - Effectiveness of Preventive Care

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Location	Provider	Adult BMI	Breast Cancer Screening	Cervical Cancer Screening	Chlamydia Screening	Childhood Immuni- zations	Colorectal Cancer Screening	Lead Screening in Children	Child BMI
SETMA 1	Aziz	95.8%	46.8%	34.6%			96.7%		
	Duncan	99.1%	51.1%	46.3%			97.9%		
	Henderson	100.0%	58.1%	46.8%			98.2%		
	Murphy	98.3%	50.2%	32.0%			97.1%		
	Thomas	98.6%	46.2%	31.8%			95.4%		
	SETMA 1 Totals:	98.7%	51.8%	41.3%			97.4%		
SETMA 2	Anthony	100.0%	56.5%	28.5%			97.3%		
	Anwar	99.8%	65.8%	29.0%			97.1%		
	Cricchio, M	100.0%	56.5%	20.1%			98.0%		
	Holly	100.0%	60.3%	34.6%			100.0%		
	Leifeste	99.8%	71.7%	38.3%			99.8%		
	Wheeler	99.8%	46.5%	50.2%			97.9%		
SETMA 2 Totals: 99		99.9%	59.8%	35.5%			98.0%		
SETMA West	Curry	97.3%	58.8%	48.5%			97.4%		
	Deiparine	99.1%	59.5%	42.0%			96.9%		
	Halbert	97.9%	52.6%	24.6%			96.1%		
	Horn	99.4%	64.0%	38.8%			99.5%		
	Qureshi	100.0%	55.6%	41.9%			100.0%		
	Satterwhite	99.7%	58.7%	32.4%			98.7%		
	Vardiman	100.0%	64.4%	33.3%			97.2%		
	Young	100.0%	74.5%	19.1%			95.0%		
SETMA West Totals: 98.9%		61.0%	36.1%			98.0%			
SETMA Totals:		99.2%	57.9%	37.4%			97.8%		

Adult Body Mass Assessment - Calculation of Body Mass Index (BMI) for patients 18 to 74 years of age.

Breast Cancer Screening - Yearly mammography screening for women 40 to 69 years of age.

Cervical Cancer Screening - Cervical cancer screening for women 21 to 64 years of age at least every two years.



HEDIS - Effectiveness of Preventive Care

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Chlamydia Screening - Sexually active females between the ages of 16 and 24 should have a Chlamydia screening at least yearly.

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Colorectal Cancer Screening - Patients 50 to 80 years of age should have at least one of the following: a colonoscopy within the last nine years, a flexible sigmoidoscopy within the last four years, a double contrast barium enema within the last four years or a fecal occult blood test within the last year.

Lead Screening in Children - Children should have a serum lead screening by the time of their second birthday.

Childhood Body Mass Assessment - Children 2 to 18 years of age should have the Body Mass Index (BMI) percentile documented.