



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2011 through Dec 31, 2011

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	--
	Duncan	--	22.2%
	Halbert	--	--
	Henderson	25.0%	38.7%
	Murphy	--	0.0%
	Palang	--	--
	Thomas	--	0.0%
SETMA 1 Totals:		25.0%	31.4%
SETMA 2	Abbas	--	--
	Anthony	14.8%	33.3%
	Anwar	--	--
	Colbert	12.9%	60.8%
	Cricchio, A	--	100.0%
	Cricchio, M	0.0%	59.5%
	Deiparine	--	--
	Holly	--	--
	Leifeste	--	--
	Murphy	--	--
	Wheeler	11.3%	67.3%
SETMA 2 Totals:		12.9%	57.0%
SETMA West	Curry	0.0%	37.5%
	Deiparine	--	6.7%
	Halbert	--	--
	Horn	--	78.6%
	Qureshi	--	69.2%
	Satterwhite	0.0%	0.0%



NQF - Pediatric Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2011 through Dec 31, 2011

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA West	Vardiman	--	20.0%
SETMA West Totals:		0.0%	43.5%
SETMA Totals:		13.0%	53.0%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.