

HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only Encounter Date(s): Jan 1, 2012 through Dec 31, 2012

| Location | Provider | Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis | Appropriate Testing for Children with Pharyngitis | Appropriate Treatment for Children with Upper Respiratory Infection |
|--------------------------|-----------|--|---|--|
| SETMA 1 | Aziz | 50.0% | 100.0% | |
| | Deiparine | | | |
| | Duncan | 11.1% | 54.5% | 25.0% |
| | Halbert | | | |
| | Henderson | 100.0% | 21.1% | 10.5% |
| | Holly | | | |
| | Murphy | | 25.0% | 0.0% |
| | Palang | 0.0% | | |
| | Thomas | | | |
| SETMA 1 Totals: | | 16.0% | 34.3% | 15.6% |
| SETMA 2 | Anthony | 100.0% | 21.0% | 0.0% |
| | Anwar | 100.0% | | 0.0% |
| | Cash | 0.0% | 0.0% | |
| | Colbert | | 58.7% | 0.0% |
| | Holly | | | |
| | LaBorde | | | |
| | Leifeste | | 0.0% | 100.0% |
| | Read | 100.0% | 40.0% | |
| | Wheeler | 60.0% | 62.3% | 11.1% |
| SETMA 2 Totals: | | 66.7% | 48.9% | 7.4% |
| SETMA Mid County | Castro | | | |
| | George | 75.0% | 100.0% | |
| | Read | | | |
| | Shepherd | 100.0% | 0.0% | |
| | Thomas | 50.0% | | |
| SETMA Mid County Totals: | | 61.5% | 25.0% | |



HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only Encounter Date(s): Jan 1, 2012 through Dec 31, 2012

| Location | Provider | Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis | Appropriate Testing for Children with Pharyngitis | Appropriate Treatment for Children with Upper Respiratory Infection |
|--------------------|-----------|--|---|--|
| SETMA West | Darden | 33.3% | 31.6% | 0.0% |
| | Deiparine | | 0.0% | |
| | Halbert | | 0.0% | |
| | Horn | 0.0% | 11.1% | 0.0% |
| | Qureshi | 0.0% | 66.7% | 50.0% |
| | Vardiman | 67.9% | 33.3% | 100.0% |
| SETMA West Totals: | | 43.5% | 24.4% | 10.0% |
| SETMA Totals: | | 40.9% | 45.3% | 11.4% |

Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis - Patients 18 to 64 years of age who were diagnosed with acute bronchitis and not prescribed an antibiotic.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.

Appropriate Treatment for Children with Upper Respiratory Infection - Patients 3 months to 18 years of age who were diagnosed with upper respiratory infection and not prescribed an antibiotic.