

NQF - Care for Older Adults

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2012 Through Dec 30, 2012

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA 1	Aziz	53.4%	67.4%	98.5%	95.8%
	Deiparine	0.0%	55.7%	87.0%	95.1%
	Duncan	87.4%	67.6%	96.4%	99.4%
	Halbert	0.0%	100.0%	0.0%	0.0%
	Henderson	64.8%	66.7%	99.8%	99.8%
	Holly	98.9%	72.1%	100.0%	98.6%
	Murphy	60.1%	62.8%	98.2%	99.3%
	Palang	31.3%	52.8%	99.1%	99.5%
	Thomas	52.8%	33.1%	100.0%	97.2%
SETMA 1 Totals:		57.3%	62.8%	98.4%	98.6%
SETMA 2	Anthony	99.0%	71.1%	99.6%	99.9%
	Anwar	99.1%	79.0%	99.6%	98.6%
	Cash	89.8%	58.4%	85.4%	94.8%
	Colbert				
	Holly	100.0%	74.1%	100.0%	100.0%
	LaBorde	98.1%	74.6%	96.8%	99.4%
	Leifeste	92.0%	85.3%	99.0%	98.8%
	Read	97.9%	72.1%	99.8%	98.9%
	Wheeler	99.0%	73.4%	99.6%	99.5%
SETMA 2 Totals:		96.9%	74.8%	98.2%	98.7%
SETMA Mid County	Castro	94.4%	19.2%	99.3%	93.1%
	George	38.6%	38.7%	100.0%	96.7%
	Read	94.0%	49.2%	98.0%	96.0%
	Shepherd	20.3%	50.4%	99.8%	88.5%
	Thomas	50.2%	27.1%	96.9%	94.7%
SETMA Mid County Totals:		39.4%	39.1%	98.8%	92.5%
SETMA West	Darden	55.1%	65.8%	98.8%	99.8%



NQF - Care for Older Adults

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2012 Through Dec 30, 2012

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA West	Deiparine	55.2%	62.3%	97.7%	98.0%
	Halbert	43.9%	52.3%	95.5%	92.4%
	Horn	90.7%	67.4%	99.5%	99.0%
	Qureshi	42.8%	59.1%	98.1%	97.8%
	Vardiman	70.0%	55.1%	93.7%	94.8%
	SETMA West Totals:	57.9%	60.0%	97.2%	96.2%
SETMA Totals:		66.7%	62.9%	98.1%	97.3%

Counseling For Physical Activity - Patients 65 and older should be counseled to engage in physical activity.

Colorectal Cancer Screening - Patients 50 to 80 years of age should have at least one of the following: a colonoscopy within the last nine years, a flexible sigmoidoscopy within the last four years, a double contrast barium enema within the last four years or a fecal occult blood test within the last year.

Fall Risk Assessment - Patients 65 and older should have a fall risk assessment completed at least yearly.

Urinary Incontinence Assessment - Patients 65 and older should have urinary incontinence assessed at least every six months.