



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2015 through Dec 31, 2015

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	--
	Duncan	--	100.0%
	Henderson	0.0%	16.7%
	Holly	--	--
	Le	0.0%	50.0%
	Murphy	--	--
	Palang	--	--
	Thomas	--	--
SETMA 1 Totals:		0.0%	33.3%
SETMA 2	Akhter	--	--
	Anthony	1.7%	33.3%
	Anwar	--	--
	Cash	--	--
	Colbert	2.0%	50.8%
	Cricchio, M	--	--
	Holly	--	--
	Kansara	--	--
	Khan	--	--
	Smith	0.0%	100.0%
	Wheeler	0.0%	72.7%
SETMA 2 Totals:		1.5%	57.9%
SETMA Lumberton	Anwar	--	--
	Dao	0.0%	66.7%
	Elliott	--	--
	Foster	1.4%	47.7%
	Holly	--	--



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2015 through Dec 31, 2015

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA Lumberton	Le	0.0%	0.0%
	Minsloff	0.0%	12.5%
	Read	--	0.0%
SETMA Lumberton Totals:		0.4%	40.0%
SETMA Mid County	Arcala, A	--	--
	Arcala, F	--	--
	Castro	--	--
	Cricchio, M	--	--
	George	--	--
	Shepherd	--	--
	Thomas	--	--
SETMA Mid County Totals:		--	--
SETMA Orange	Castro	--	--
	Cox	--	0.0%
	Green	--	46.2%
	McClure	--	--
	Read	--	0.0%
	Shepherd	--	--
SETMA Orange Totals:		--	37.5%
SETMA PA	Arcala, A	--	--
	Arcala, F	--	--
	Castro	--	--
	Shepherd	--	--
SETMA PA Totals:		--	--
SETMA West	Darden	--	0.0%
	Deiparine, C	--	--



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2015 through Dec 31, 2015

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA West	Halbert	--	--
	Horn	0.0%	76.9%
	Khan	--	--
	Qureshi	--	--
SETMA West Totals:		0.0%	66.7%
SETMA Totals:		1.3%	49.8%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.