



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2016 through Dec 31, 2016

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	100.0%
	Cash	--	--
	Duncan	--	83.3%
	Henderson	0.0%	30.4%
	Holly	--	100.0%
	Le	0.0%	43.8%
	Murphy	--	--
	Palang	--	--
	Thomas	--	--
SETMA 1 Totals:		0.0%	44.7%
SETMA 2	Akhter	--	--
	Anthony	0.4%	45.5%
	Anwar	--	--
	Cash	--	--
	Colbert	0.5%	53.2%
	Kansara	--	--
	Lampkin	--	--
	Smith	1.6%	57.7%
	Wheeler	0.0%	72.9%
SETMA 2 Totals:		0.6%	60.1%
SETMA Lumberton	Dao	0.0%	63.2%
	Elliott	--	0.0%
	Foster	0.0%	25.0%
	Minsloff	0.0%	81.1%
	Read	--	14.3%
SETMA Lumberton Totals:		0.0%	69.3%



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2016 through Dec 31, 2016

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA Mid County	Akhter	--	--
	Arcala, A	--	100.0%
	Castro	--	--
	Cox	--	--
	Cricchio, M	--	0.0%
	Feldschau	--	33.3%
	George	--	--
	Green	--	100.0%
	Henderson	--	--
	Le	0.0%	--
	Ramirez	--	--
	Read	--	--
	Shepherd	--	0.0%
	Thomas	--	--
SETMA Mid County Totals:		0.0%	42.9%
SETMA Orange	Castro	--	--
	Cox	--	50.0%
	Green	--	71.4%
	Read	--	50.0%
	Shepherd	--	--
SETMA Orange Totals:		--	63.6%
SETMA PA	Arcala, A	--	--
SETMA PA Totals:		--	--
SETMA West	Darden	--	80.0%
	Deiparine, C	--	--
	Foster	--	--



NQF - Pediatric Measures

E & M Codes: Clinic Only
Encounter Date(s): Jan 1, 2016 through Dec 31, 2016

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA West	Halbert	--	0.0%
	Horn	0.0%	58.8%
	Khan	--	--
	Murphy	--	--
	Qureshi	--	--
SETMA West Totals:		0.0%	60.9%
SETMA Totals:		0.4%	63.9%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.