



NQF - Pediatric Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2017 through Dec 31, 2017

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	--
	Duncan	--	80.0%
	Halbert	--	--
	Henderson	0.0%	42.9%
	Holly	--	--
	Le	0.0%	15.8%
	Murphy	--	100.0%
	Palang	--	--
	Qureshi	--	--
	Ramirez	--	--
	Smith	--	--
	Thomas	--	--
SETMA 1 Totals:		0.0%	37.0%
SETMA 2	Akhter	--	--
	Anthony	0.0%	30.0%
	Anwar	--	--
	Cash	--	--
	Colbert	0.0%	65.8%
	Darden	--	--
	Kansara	--	--
	Le	--	--
	Murphy	--	--
	Palang	--	--
	Smith	0.0%	54.1%
	Wheeler	0.0%	83.0%
	SETMA 2 Totals:		0.0%



NQF - Pediatric Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2017 through Dec 31, 2017

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA Lumberton	Castro	--	100.0%
	Colbert	0.0%	75.4%
	Feldschau	--	--
	Foster	0.0%	76.0%
	Halbert	--	--
	Holly	--	--
	Horn	--	--
	Kansara	--	--
	Khan	--	--
	Palang	--	--
	Qureshi	--	--
	Ramirez	--	62.5%
	SETMA Lumberton Totals:		0.0%
SETMA Mid County	Castro	--	--
	Feldschau	--	25.0%
	George	--	--
	Le	0.0%	22.2%
	Ramirez	--	--
	Shepherd	--	--
	Thomas	--	--
SETMA Mid County Totals:		0.0%	23.1%
SETMA Orange	Castro	--	--
	Cox	--	50.0%
	Green	--	60.0%
	Shepherd	--	--
SETMA Orange Totals:		--	57.9%



NQF - Pediatric Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2017 through Dec 31, 2017

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA West	Akhter	--	--
	Darden	--	0.0%
	Deiparine, C	--	--
	Halbert	--	25.0%
	Horn	0.0%	75.0%
	Khan	--	--
	Qureshi	--	--
SETMA West Totals:		0.0%	50.0%
SETMA Totals:		0.0%	64.5%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.